

Manuscript Number:

Title: Mast cells and inflammatory bowel disease

Article Type: 25: Gastrointestinal (2015)

Corresponding Author: Prof. Dr. Guy Boeckxstaens,

Corresponding Author's Institution:

First Author: Guy Boeckxstaens

Order of Authors: Guy Boeckxstaens

Abstract: Inflammatory bowel diseases (IBD), consisting of ulcerative colitis (UC) and Crohn's disease (CD), are chronic immune-mediated diseases of the gut. Here, the potential role of mast cells (MC) is discussed, mainly focusing on preclinical studies. MC can be activated by antigen-mediated crosslinking of immunoglobulin receptors, by free light chains of immunoglobulins, stress and ATP. Upon activation, MC release bioactive mediators, of which the serine proteases mMCP-6 and Prss31 were shown to be involved in the development of acute colitis. Inhibition of MCs by activation of the inhibitory receptor LIMR3 or inhibitors of proteases may therefore represent new therapeutic targets to treat IBD. Human data are however lacking.

Suggested Reviewers:

**\*Conflict of interest statement**

Conflicts of interest:

Nothing to declare

### **Highlights**

- **ATP activates MC via P2X7 receptors leading to colitis**
- **Activation of the inhibitory receptor LMR3 on MC reduces colitis**
- **The MC proteases MCP-6 and Prss31 are involved in DSS and TNBS colitis**
- **Protease inhibitors or LMR3 agonists could be interesting compounds to treat IBD**

# **Mast cells and inflammatory bowel disease**

Guy Boeckxstaens

Address of correspondence:

Guy E. Boeckxstaens

Div. Gastroenterology

Translational Research Center for Gastrointestinal Disorders (TARGID)

KULeuven, University Hospital Leuven,

Herestraat 49

3000 Leuven

Belgium

Email: [guy.boeckxstaens@med.kuleuven.be](mailto:guy.boeckxstaens@med.kuleuven.be)

### **Abstract (100-120 words)**

Inflammatory bowel diseases (IBD), consisting of ulcerative colitis (UC) and Crohn's disease (CD), are chronic immune-mediated diseases of the gut. Here, the potential role of mast cells (MC) is discussed, mainly focusing on preclinical studies. MC can be activated by antigen-mediated crosslinking of immunoglobulin receptors, by free light chains of immunoglobulins, stress and ATP. Upon activation, MC release bioactive mediators, of which the serine proteases mMCP-6 and Prss31 were shown to be involved in the development of acute colitis. Inhibition of MCs by activation of the inhibitory receptor LMR3 or inhibitors of proteases may therefore represent new therapeutic targets to treat IBD. Human data are however lacking.

### **Highlights**

- **ATP activates MC via P2X7 receptors leading to colitis**
- **Activation of the inhibitory receptor LMR3 on MC reduces colitis**
- **The MC proteases MCP-6 and Prss31 are involved in DSS and TNBS colitis**
- **Protease inhibitors or LMR3 agonists could be interesting compounds to treat IBD**

## **Introduction**

Crohn's disease (CD) [1] and ulcerative colitis (UC) [2] are the two main types of idiopathic chronic inflammatory disorders (IBD) of the gastrointestinal tract, both associated with a significant impact on the quality of life of the affected individuals. While CD can affect any part of the gut, albeit with a predilection for the terminal ileum, UC rather typically starts in the rectum and may extend all the way up to the caecal area. Despite extensive scientific effort, the exact underlying cause of IBD remains unclear. It is however becoming increasingly clear that a dysregulated mucosal response to the commensal gut flora is involved, in particular in genetically susceptible individuals. In UC, this immune response involves an atypical Th2 response mediated by non-classic natural killer T cells. In contrast, in Crohn's disease, imbalance between predominantly Th1 or Th17 cells and naturally regulatory T cells is proposed to underlie the mucosal inflammation and associated damage. In the present review, the evidence, albeit scarce, supporting a potential contribution of mast cells will be summarized. Mast cells are particularly interesting as they are not only involved in the immediate and delayed defense against foreign antigens [3], but also release a mixture of mediators known to affect mucosal barrier function[4-6], thereby increasing the access of intraluminal antigens to the mucosal immune system.

## **Mast cells: general aspects**

Mast cells arise from hematopoietic progenitors in the bone marrow that in contrast to most other immune cells only mature when they arrived at their final destination, i.e. all vascularized tissues but especially the skin and mucosal tissues. Typically, mast cells reside in the mucosa at anatomical sites directly exposed to the environment, such as the skin, the airways and the gastrointestinal tract, where they are ideally positioned to respond to foreign antigens that have crossed the epithelial barrier[6]. Their number can vary significantly depending on the tissue studied but also during infections or

inflammation. Both in man and mice, two main types of mast cells can be identified, largely based on location and protease content. In mice, connective tissue mast cells and mucosal mast cells are distinguished. Connective tissue mast cells reside mainly around venules[7] and nerve fibers and in serosal cavities. Mucosal mast cells on the other hand are present in the mucosa of the gut and lung, but in low numbers[8]. In humans, mast cells are characterized by the proteases they contain: the MC<sub>T</sub>-type mast cells containing tryptase but little or no chymase, and the MC<sub>TC</sub>-type mast cells containing tryptase and chymase[9]. Almost all human mucosal mast cells are MC<sub>T</sub> while MC<sub>TC</sub> are typically located in the lamina propria and submucosa.

Mast cells can be activated by a wide variety of stimuli. The most well known mechanism of activation is cross-linking of the high affinity IgE receptor FcεRI. This receptor binds antigen-specific IgE antibodies captured from the circulation by mast cell processes extended across the vessel wall [7]. Binding of these antibodies with their respective antigen leads to crosslinking and subsequently aggregation of FcεRIs thereby activating downstream signaling pathways and triggering the release of biologically active mediators [10]. In addition, mast cells can be activated via toll like receptors, G-protein coupled receptors, various complement peptides and platelet activating factor, illustrating its crucial role in immune defense [3]. Of note, mast cells are armed with granules containing preformed and stored mediators, such as vasoactive amines like histamine, and proteases such as tryptases and chymases, enabling an immediate response. In addition, mast cells can synthesize and release lipid mediators such as prostaglandins and leukotrienes within 1-2 hours after activation. Mast cells also release cytokines and chemokines, a process that typically occurs later and is associated with more chronic responses. The wide range of mediators released undoubtedly underlie the many different functions of mast cells and their involvement in a diversity of mechanisms, including secretion and epithelial permeability, immune cell recruitment and activation, blood flow, coagulation and vascular permeability, wound healing and

fibrosis and neuro-immune interaction altering peristalsis, bronchoconstriction and pain [11].

### **Evidence in humans supporting a role for mast cells in IBD**

To date, only few studies have evaluated the potential involvement of mast cells in IBD, and most of them date back many years ago and are rather descriptive reporting on the number of mast cells. Gelbmann et al. [12] for example describe accumulation of mast cells particularly in hypertrophied and fibrotic muscularis propria in strictures of CD patients, suggesting an important role in the process of fibrosis. Others report on increased mast cells in the mucosa, submucosa, muscularis propria and surrounding fat in active CD [13]. More recently, increased mast cell numbers were counted in the colon of both CD and UC patients compared to controls [14] [15]. Activation of mast cells on the other hand is suggested by increased expression or release of mast cell mediators in the mucosa of IBD patients [16] [17] or by demonstration of increased levels of the N-methylhistamine, one of the metabolites of histamine, in the urine of IBD patients compared to patients in remission and controls [18]. Although the above-mentioned observations may suggest a role for mast cells in IBD, this evidence is rather indirect and could also be secondary to the inflammatory response in the mucosa. The ultimate proof, i.e. clinical improvement by treatment with mast cell stabilizers or antagonist blocking the biological effect of mast cell mediators, is however currently lacking.

### **Mast cells and colitis: preclinical data**

Several models of colitis are routinely used that can largely be divided in models of acute colitis[19], mostly due to exposure of the intestine to compounds damaging the intestinal barrier (dextran sulphate sodium (DSS)) in drinking water, enema of trinitrobenzene dissolved in alcohol) or more chronic models such as T cell transfer colitis or delayed-type hypersensitivity colitis (TNBS, dinitrofluoro-benzene). Clearly,



the role of mast cells in colitis will largely depend on the type of model studied. Below, recent new data supporting a role for mast cells in colitis will be briefly summarized according to the type of colitis.

### ***Acute colitis***

Previous studies have yielded conflicting data regarding the role of mast cells in acute colitis [20]. Kurashima et al. however nicely demonstrated a pivotal role for ATP-mediated activation of mast cells in DSS and TNBS colitis [15]. These authors demonstrated that treatment of mice with an antibody against the ATP receptor P2X7 inhibits mast cell activation and subsequent intestinal inflammation. Moreover, mast cells expressing P2X7 purinoceptors in colons of mice with colitis and of patients with CD are increased in number. The most compelling evidence however is revealed by reconstitution experiments in mast cell deficient *Kit<sup>W-sh/W-sh</sup>* mice. Intestinal inflammation triggered by DSS or TNBS in these mutant mice is ameliorated while reconstitution with wild type but not P2X7<sup>-/-</sup> mast cells results in susceptibility to develop colitis in response to DSS or TNBS [15]. Moreover, the authors showed increased ATP production in intestinal inflammation and demonstrated that ATP triggers MC activation through P2X7 purinoreceptors. Taken together, these data indicate that tissue damage induced by DSS or TNBS leads to ATP release subsequently activating MC by interacting with P2X7 receptors. These data reveal a role for MC in both the initiation and exacerbation of intestinal inflammation [15]. Of interest, the same group very recently showed that this pathway can be blocked by activation of the inhibitory receptor CD300f (also called leucocyte mono-immunoglobulin-like receptor 3 or LMIR3) on mast cells by its ligand ceramide [21]. Mice deficient for LMIR3 developed more severe DSS colitis. Moreover, reconstitution of mast cell deficient mice with LMIR3<sup>-/-</sup> MC exhibited more severe colitis than mice that received wild type MC. Previously, the authors have shown that ceramide is a ligand of LMIR3 leading to

inhibition of IgE-dependent MC activation [22]. In their recent paper, they take this finding further and show that ceramide-LIMR3 interaction not only inhibits ATP-induced MC activation but also suppresses DSS colitis. These data are of great interest indicating LIMR3-targeted compounds such as ceramide as a novel therapeutic strategy for IBD [21].

In mice, mast cells store a variety of proteases on their secretory granules. These enzymes are of particular interest as they are involved in proteolytic processing of other molecules but can also act as signaling molecules by cleavage and activation of Protease-activated receptors (PARs) [23]. The latter are abundantly expressed and trigger a wide array of pro-inflammatory processes upon activation, including colitis[24]. Two recent papers have studied the role of mast cell serine proteases in DSS colitis, in particular mouse MC protease-6 (mMCP-6), mMCP-7 and protease serine member S31 (Prss31) [25,26]. Hamilton et al. [25] elegantly showed that mMCP-6 null mice exposed to DSS had significantly less weight loss as well as lower pathological and endoscopical disease scores compared to mMCP-6 expressing mice. In line, the levels of the pro-inflammatory cytokines IL-6 and IL-1 $\beta$  and the chemokines CXCL1 and CXCL2, known to attract neutrophils, were significantly reduced in the mMCP-6 null mice. Of interest, similar findings were obtained in the TNBS colitis model. In a subsequent study, the same group showed that also Prss31 null mice lost less weight, had lower histopathological scores and contained less CXCL2 and IL-6 mRNA in their colons compared to WT mice following treatment with DSS. Taken together, these data provide strong evidence that both mMCP-6 and Prss31 contribute to the development of colitis and thus may represent interesting new targets for treatment of IBD [26].

### **Delayed-type hypersensitivity colitis**

After sensitization, re-exposure to oxazolone, TNBS or dinitrofluorobenzene leads to a delayed-type of colitis. Classically, MC are activated by antigen mediated crosslinking of

membrane bound IgE antibodies. Recently, Hoving et al. [27] provided evidence for IgE-mediated activation of MC in oxazolone colitis. In vivo neutralization of IgE protected mice from oxazolone-induced colitis, a finding associated with a reduction in mast cell numbers and a reduction in the MC degranulation product MMCP-1 in serum. Moreover, mice that are unable to produce IgE were also protected against oxazolone-induced colitis. Next, the authors showed that IgE antibodies were produced by B cells in response to IL-13 released by CD4<sup>+</sup> Th2 cells. Along the same line, Rijnierse et al. provided evidence that mast activation by immunoglobulin-free light chains (IgLCs) of immunoglobulins are involved in DNBS colitis [28]. Treatment of mice with the IgLC antagonist F991 prevented MC activation and abrogated the development of diarrhea, cellular infiltration and colonic lymphoid follicle hyperplasia. Passive immunization with antigen specific IgLCs and subsequent rectal challenge with the hapten elicited MC activation and increased vascular permeability in the colon of mice. Of note, serum concentrations of IgLCs in CD was increased while more IgLCs were detected in the ileum and colon of patients with IBD [28]. Based on the above, IgE dependent and IgE-independent MC activation may be involved in IBD. Further data in human are however awaited.

### **Stress, mast cells and IBD**

A recent large prospective population-based study in 704 IBD patients identified perceived stress, negative affect or mood and major life events to be associated with symptomatic flares [29]. Knowing that stress is an important trigger of MC activation [30], MC could theoretically be involved in the triggering of these flares, perhaps by increasing permeability thereby exposing the immune system to increased levels of intraluminal antigens. Increased permeability has indeed been reported using biopsies of IBD patients [14] [31], whereas reactivation of colitis by stress was reported already many years ago. This effect was dependent on CD4<sup>+</sup> T cells and increased permeability

[32]. In 2009, evidence was reported suggesting that chronic psychological stress, through the release of substance P, triggers the expression and release of corticotrophin releasing hormone by eosinophils. This stress hormone then subsequently activates MC leading to epithelial barrier dysfunction [33]. The above-mentioned studies would suggest that stress-induced flares in IBD may at least partly be explained by MC-mediated changes in barrier function and subsequent reactivation of the immune response.

### **Summary and conclusion**

From the studies summarized above, MC seem to be involved both in the acute and more sustained inflammatory response, at least in preclinical models of IBD. As shown in Figure 1, MC can be activated by stress via eosinophil-derived CRH, IgLC, IgE or ATP leading to the release of a wide variety of mediators, including proteases, histamine, chemokines and cytokines, leading to the attraction of inflammatory cells, changes in barrier function, tissue remodeling etc. Increased permeability may contribute to increased influx of intraluminal antigens either activating MC or CD4+ T cells further contributing to the inflammatory response. This process can be antagonized by inhibition of MC activation via ceramide binding to the inhibitory receptor LMR3. To what extent this also applies to IBD patients and to what extent MC are indeed a target for treatment remains to be proven. Clearly, the most convincing evidence will have to come from clinical studies evaluating new compounds in patients, however, data from randomized clinical trials are currently available. Nevertheless, based on the data summarized here, protease inhibitors[23,34] and ceramide[21] may represent interesting targets hopefully improving treatment of IBD patients.

## Acknowledgements

G. Boeckxstaens is supported by a governmental (Odysseus programme, G.0905.07) of the Research Foundation Flanders (FWO) and university grant (Global Opportunities for Associations GOA 14.011). GB is also supported by an European Research Council (ERC) Advanced research grant (Grant Agreement Number 340101).

## Legend to Figure 1

Schematic representation of the possible involvement of MC in IBD. For more detailed information, see main text.

1. Baumgart DC, Sandborn WJ: **Crohn's disease**. *Lancet* 2012, **380**:1590-1605.
2. Ordas I, Eckmann L, Talamini M, Baumgart DC, Sandborn WJ: **Ulcerative colitis**. *Lancet* 2012, **380**:1606-1619.
3. Abraham SN, St John AL: **Mast cell-orchestrated immunity to pathogens**. *Nat Rev Immunol* 2010, **10**:440-452.
4. Sayed BA, Brown MA: **Mast cells as modulators of T-cell responses**. *Immunol Rev* 2007, **217**:53-64.
5. Bischoff SC: **Physiological and pathophysiological functions of intestinal mast cells**. *Semin Immunopathol* 2009, **31**:185-205.
6. Reber LL, Sibilano R, Mukai K, Galli SJ: **Potential effector and immunoregulatory functions of mast cells in mucosal immunity**. *Mucosal Immunol* 2015, **8**:444-463.
7. Cheng LE, Hartmann K, Roers A, Krummel MF, Locksley RM: **Perivascular mast cells dynamically probe cutaneous blood vessels to capture immunoglobulin E**. *Immunity* 2013, **38**:166-175.
8. Gurish MF, Austen KF: **Developmental origin and functional specialization of mast cell subsets**. *Immunity* 2012, **37**:25-33.
9. Weidner N, Austen KF: **Heterogeneity of mast cells at multiple body sites. Fluorescent determination of avidin binding and immunofluorescent determination of chymase, tryptase, and carboxypeptidase content**. *Pathol Res Pract* 1993, **189**:156-162.
10. Gilfillan AM, Tkaczyk C: **Integrated signalling pathways for mast-cell activation**. *Nat Rev Immunol* 2006, **6**:218-230.
11. Bischoff SC, Kramer S: **Human mast cells, bacteria, and intestinal immunity**. *Immunol Rev* 2007, **217**:329-337.
12. Gelbmann CM, Mestermann S, Gross V, Kollinger M, Scholmerich J, Falk W: **Strictures in Crohn's disease are characterised by an accumulation**

**of mast cells colocalised with laminin but not with fibronectin or vitronectin.** *Gut* 1999, **45**:210-217.

13. Andoh A, Deguchi Y, Inatomi O, Yagi Y, Bamba S, Tsujikawa T, Fujiyama Y: **Immunohistochemical study of chymase-positive mast cells in inflammatory bowel disease.** *Oncol Rep* 2006, **16**:103-107.

14•. Vivinus-Nebot M, Frin-Mathy G, Bziouche H, Dainese R, Bernard G, Anty R, Filippi J, Saint-Paul MC, Tulic MK, Verhasselt V, et al.: **Functional bowel symptoms in quiescent inflammatory bowel diseases: role of epithelial barrier disruption and low-grade inflammation.** *Gut* 2014, **63**:744-752.

Interesting study showing that patients with IBD have increased mucosal permeability compared to healthy controls.

15••. Kurashima Y, Amiya T, Nochi T, Fujisawa K, Haraguchi T, Iba H, Tsutsui H, Sato S, Nakajima S, Iijima H, et al.: **Extracellular ATP mediates mast cell-dependent intestinal inflammation through P2X7 purinoceptors.** *Nat Commun* 2012, **3**:1034.

This study provides compelling evidence that ATP release is increased in intestinal inflammation and activates MC through P2X7 receptors

16. Raithel M, Schneider HT, Hahn EG: **Effect of substance P on histamine secretion from gut mucosa in inflammatory bowel disease.** *Scand J Gastroenterol* 1999, **34**:496-503.

17. Raithel M, Winterkamp S, Pacurar A, Ulrich P, Hochberger J, Hahn EG: **Release of mast cell tryptase from human colorectal mucosa in inflammatory bowel disease.** *Scand J Gastroenterol* 2001, **36**:174-179.

18. Winterkamp S, Weidenhiller M, Otte P, Stolper J, Schwab D, Hahn EG, Raithel M: **Urinary excretion of N-methylhistamine as a marker of disease activity in inflammatory bowel disease.** *Am J Gastroenterol* 2002, **97**:3071-3077.

19. te Velde AA, de Kort F, Sterrenburg E, Pronk I, ten Kate FJ, Hommes DW, van Deventer SJ: **Comparative analysis of colonic gene expression of three experimental colitis models mimicking inflammatory bowel disease.** *Inflamm Bowel Dis* 2007, **13**:325-330.

20. Rijnerse A, Nijkamp FP, Kraneveld AD: **Mast cells and nerves tickle in the tummy: implications for inflammatory bowel disease and irritable bowel syndrome.** *Pharmacol Ther* 2007, **116**:207-235.

21••. Matsukawa T, Izawa K, Isobe M, Takahashi M, Maehara A, Yamanishi Y, Kaitani A, Okumura K, Teshima T, Kitamura T, et al.: **Ceramide-CD300f binding suppresses experimental colitis by inhibiting ATP-mediated mast cell activation.** *Gut* 2015.

This study reports on the role of the inhibitory receptor LMIR3 which inhibits MC activation and reduces colitis. Ceramide, an activator of this receptor, also reduces colitis, providing evidence for a new therapeutic approach for IBD.

22. Nakahashi-Oda C, Tahara-Hanaoka S, Shoji M, Okoshi Y, Nakano-Yokomizo T, Ohkohchi N, Yasui T, Kikutani H, Honda S, Shibuya K, et al.: **Apoptotic cells suppress mast cell inflammatory responses via the CD300a immunoreceptor.** *J Exp Med* 2012, **209**:1493-1503.

23••. Vergnolle N: **Protease inhibition as new therapeutic strategy for GI disease.** *Gut* 2015, in press.

Comprehensive review on the proteases as new target to treat intestinal diseases, including IBD.

24. Vergnolle N, Cellars L, Mencarelli A, Rizzo G, Swaminathan S, Beck P, Steinhoff M, Andrade-Gordon P, Bunnett NW, Hollenberg MD, et al.: **A role for proteinase-activated receptor-1 in inflammatory bowel diseases.** *J Clin Invest* 2004, **114**:1444-1456.
  25. Hamilton MJ, Sinnamon MJ, Lyng GD, Glickman JN, Wang X, Xing W, Krilis SA, Blumberg RS, Adachi R, Lee DM, et al.: **Essential role for mast cell tryptase in acute experimental colitis.** *Proc Natl Acad Sci U S A* 2011, **108**:290-295.
  - 26•. Hansbro PM, Hamilton MJ, Fricker M, Gellatly SL, Jarnicki AG, Zheng D, Frei SM, Wong GW, Hamadi S, Zhou S, et al.: **Importance of mast cell Prss31/transmembrane tryptase/tryptase-gamma in lung function and experimental chronic obstructive pulmonary disease and colitis.** *J Biol Chem* 2014, **289**:18214-18227.
- Interesting study identifying the protease Prss31 as mediator released by MC and contributing to the development of colitis in mice.
27. Hoving JC, Kirstein F, Nieuwenhuizen NE, Fick LC, Hobeika E, Reth M, Brombacher F: **B cells that produce immunoglobulin E mediate colitis in BALB/c mice.** *Gastroenterology* 2012, **142**:96-108.
  28. Rijnierse A, Redegeld FA, Blokhuis BR, Van der Heijden MW, Te Velde AA, Pronk I, Hommes DW, Nijkamp FP, Koster AS, Kraneveld AD: **Ig-free light chains play a crucial role in murine mast cell-dependent colitis and are associated with human inflammatory bowel diseases.** *J Immunol* 2010, **185**:653-659.
  29. Bernstein CN, Singh S, Graff LA, Walker JR, Miller N, Cheang M: **A prospective population-based study of triggers of symptomatic flares in IBD.** *Am J Gastroenterol* 2010, **105**:1994-2002.
  30. Soderholm JD, Perdue MH: **Stress and gastrointestinal tract. II. Stress and intestinal barrier function.** *Am J Physiol Gastrointest Liver Physiol* 2001, **280**:G7-G13.
  31. Wallon C, Persborn M, Jonsson M, Wang A, Phan V, Lampinen M, Vicario M, Santos J, Sherman PM, Carlson M, et al.: **Eosinophils express muscarinic receptors and corticotropin-releasing factor to disrupt the mucosal barrier in ulcerative colitis.** *Gastroenterology* 2011, **140**:1597-1607.
  32. Qiu BS, Vallance BA, Blennerhassett PA, Collins SM: **The role of CD4+ lymphocytes in the susceptibility of mice to stress-induced reactivation of experimental colitis.** *Nat Med* 1999, **5**:1178-1182.
  33. Zheng PY, Feng BS, Oluwole C, Struiksma S, Chen X, Li P, Tang SG, Yang PC: **Psychological stress induces eosinophils to produce corticotrophin releasing hormone in the intestine.** *Gut* 2009, **58**:1473-1479.
  34. Tremaine WJ, Brzezinski A, Katz JA, Wolf DC, Fleming TJ, Mordenti J, Strenkoski-Nix LC, Kurth MC, Group AUCS: **Treatment of mildly to moderately active ulcerative colitis with a tryptase inhibitor (APC 2059): an open-label pilot study.** *Aliment Pharmacol Ther* 2002, **16**:407-413.

Figure 1

